

Spectral representation for on-shell massless double box

N.I.Ussyukina

*Institute for Nuclear Physics, Moscow State University, 119899
Moscow, Russia*

E-mail address: ussyuk@theory.npi.msu.su

A method is presented for obtaining the ε -expansion
for on-shell massless scalar double-box diagram.

1. There are several reasons why calculation of two-loop diagrams is important. The study of two-loop diagrams is required by increasing precision of experiments. Examination of multi-jet processes requires analytical calculations of two-loop diagrams. At last, consideration of such diagrams helps us to develop and apply new methods of loop calculations.

One-loop on-shell four-point diagram in ϕ^4 -theory has been considered in [1]. As compared with the one-loop case, the evaluation of two-loop diagrams is technically much more complicated and less investigated. Off-shell double-box problem in 4-dimensional space-time was solved in [2]. On-shell double-box problem is more complicated than the off-shell one because the appearance of the infrared divergences. Our consideration will be in the framework of dimensional regularization [3]. On-shell double-box diagram has a very high degree of divergences. Let $n = 4 + 2\varepsilon$ be the space-time dimension. Then on-shell double-box diagram $D(s, t, k_i^2)$ diverges as ε^{-4} . It is evident that the expansion up to the fourth degree is a very difficult problem. On-shell problem can be solved for example by using various reduction methods (the integration-by-part method [4], uniqueness method [5]). Such an approach was used in [6], but the result was too cumbersome. For this reason it remained the problem to look for such a scheme, that could lead to a more visible result. Such a method I'll consider in this paper. The talk is about so named Mellin expansion method. At first this method was used for multiloop calculations in [7]. With the help of this method I managed to find the visible solution for the on-shell double-box case.

The remainder of the paper is organized as follows. In Section 2 we list notations, some most useful formulas and the main idea of calculation. In Section 3 we'll carry out some spadework before to use the Mellin expansion method. In Section 4 using the Mellin expansion method we'll obtain the spectral representation for four-point function. In Section 5 we discuss the results.

2. In this section we list notations, some most useful formulas and the main idea of calculation.




Fig.1

We use the standard notations $s = (k_1 + k_2)^2$, $t = (k_2 + k_3)^2$, $k_i^2 = 0$.

Not to complicate formulae, we omit henceforth the factors that are powers of 2, π , i . These factors can be easily restored in the final results. Each line of diagrams carries a power-like factor that is pictured as a line with a label α , for example,

$$\frac{\alpha}{\quad} \Rightarrow \frac{1}{(p^2 - i0)^\alpha} \quad \text{in momentum space,}$$

$$\frac{\alpha}{\quad} \Rightarrow \frac{1}{(x^2 + i0)^\alpha} \quad \text{in coordinate space.}$$

We'll use constantly the formula for propagator Fourier-transformation from momentum to coordinate space:

$$\frac{1}{(k^2)^\alpha} \Rightarrow \frac{\Gamma(\mu - \alpha)}{\Gamma(\alpha)} \times \frac{1}{(x^2)^{\mu - \alpha}}.$$

The main idea of calculations consists in the possibility to reduce in the case $k_4^2 = k_3^2 = 0$ double-box diagram $D(s, t)$ to three-point diagram (Fig.2) $\Gamma(q_1^2, q_2^2, q_3^2)$:




Fig.2

where


$$q_1^2 = (k_1 + k_2 x)^2, \quad q_2^2 = (k_3 + k_4 y)^2, \quad q_3^2 = t \bar{x} \bar{y}, \quad \bar{x} = 1 - x, \quad \bar{y} = 1 - y,$$


with the help of equality

$$D(s, t) = \int_0^1 dx \int_0^1 dy \Gamma(q_1^2, q_2^2, q_3^2). \quad (1)$$


The three-point diagram is more simple object for calculation and for expansion in ϵ than four-point diagram and if we will know the ϵ -expansion for Γ , we'll obtain simply ϵ -expansion for D from this equality.


3. In this section we'll carry out some preparatory work before using the Mellin-expansion method. The reason for this will be clear later. Using the uniqueness method we present the three-point diagram (here it is written in coordinate space) in the following form;



$$= \varepsilon \frac{\Gamma(1+3\varepsilon)}{\Gamma^4(1+\varepsilon)\Gamma(1-2\varepsilon)}$$


$$,$$
(2)



$$= 2 \frac{\Gamma(2-\varepsilon)\Gamma^3(1+\varepsilon)}{\Gamma(1+2\varepsilon)}$$


$$.$$
(3)

We begin our calculation from the usual Feynman α -representation for diagram of Fig.3 in coordinate representation. Then we transform this α -representation to spectral representation for Fig. 4.




Fig. 3




Fig.4

4. In this section we obtain at first spectral representation for diagram of Fig.3, then using the Mellin expansion method we'll reduce it to spectral representation for four-point function. Spectral representation for diagram of Fig.3 has the form

$$\int_0^\infty d\xi \int_0^\infty d\zeta \frac{1}{\{\xi\zeta + \xi + \zeta\}^{2-\varepsilon} \{(y-z)^2 + (x-y)^2\xi + (x-z)^2\zeta\}^{2\varepsilon}}.$$

This expression can be presented in the form of two-fold Mellin integral [8]. After multiplying this Mellin expansion by the propagators that correspond to formula (3) and after transforming the result to momentum representation we obtain for two-loop three-point diagram of Fig.4 in momentum representation the following α -representation

$$\int \prod_i d\alpha_i \delta(1 - \sum_i \alpha_i) \frac{\alpha_1^{2\varepsilon} \alpha_2^{2\varepsilon} \alpha_3^{2-2\varepsilon} \varphi(\alpha_i)}{\{q^2 \alpha_1 \alpha_2 + q_1^2 \alpha_2 \alpha_3 + q_2^2 \alpha_1 \alpha_3\}}, \quad (4)$$

where

$$\varphi(\alpha_i) = \int_1^\infty dx \frac{1}{x^{1-\varepsilon}} \int_0^1 dz \frac{z^{2\varepsilon-1} (1-z)^{2\varepsilon-1}}{\{\alpha_1 x z + \alpha_2 z + \alpha_3\}^{1+\varepsilon}}.$$

This representation is very similar to ordinary α -representation for one-loop three-point function, but the representation for two-loop case include weight-function $\varphi(\alpha_i)$.

After transformations that take into account equality (1) we obtain for four-point function:

$$D(s, t) \Rightarrow \int_0^1 dx \int_0^1 dy \int_0^\infty d\xi \int_0^\infty d\zeta \frac{\zeta^\varepsilon \xi^{2\varepsilon}}{\{q^2 + \xi + \zeta\}^\varepsilon \{\xi\zeta + q_1^2 \xi + q_2^2 \zeta\}^2} \int_0^\infty d\tau \tau^{2\varepsilon-1} (1+\tau)^{1-3\varepsilon} \\ \times \int_\zeta^\infty dz \frac{z^{\varepsilon-1}}{\{(1+\tau)q^2 + \xi + z\}^{1+\varepsilon}}.$$

At this stage we can see why the previous transformations were done. For the form (3) we can carry out integration over y . Then

$$D(s, t) \Rightarrow \int_0^1 dx \int_0^\infty d\xi \int_0^\infty d\zeta \int_0^\infty d\tau \frac{\xi \tau^{2\varepsilon-1}}{\{1+\tau+s\zeta\}\{1+\tau+t\bar{x}\xi\}} \int_0^\zeta dz \frac{1}{\{q^2 \xi z + \xi + z\}^{1+\varepsilon}}, \quad (5)$$

and taking into account, that $q^2 = sx$ we obtain (here $L = t/s$)

$$D(s, t) \Rightarrow \frac{1}{s^{3-2\varepsilon}} J, \\ J = \varepsilon \int_0^\infty d\xi \int_0^\infty d\zeta \int_0^\infty dx \int_0^\infty d\tau \frac{\xi \tau^{2\varepsilon-1}}{\{1+\tau+L\xi\}\{1+\tau+\zeta\}} \\ \times \frac{1}{\{x\zeta + (1+\tau)(x+\xi+\zeta)\}^\varepsilon} \int_0^\zeta dz \frac{1}{\{xz + x + z + \xi\}^{1+\varepsilon}}. \quad (6)$$

This spectral representation is a convenient starting point for both analytical and numerical calculations. For instance, in a separate publication I will obtain one of the irreducible diagrams that appeared in [10], which corresponds to $L = -1$ as well as a much more cumbersome analytical result for arbitrary L as a quadratic form of Li_n -functions.

Conclusions. In this paper we have considered an approach to the calculation of on-shell two-loop scalar double-box diagram. This approach is based on the following tools: (i) the Feynman parametric representation, (ii) the uniqueness method, (iii) Mellin-Barnes contour integrals. We considered only scalar diagrams (corresponding to massless φ^3 -theory) because expressions occurring in realistic calculations can be reduced to such scalar integrals. We have obtained the analytical formula (6), that can be used for calculations in various concrete cases.

I am grateful to F.V.Tkachov for his interest in this work, useful discussions and help. The research was supported in part by the Russian Fund for Basic Research (grant 95-02-05794).

References

- [1] W.L.van Neerven. Nucl. Phys. B268 (1986) 453.
- [2] N.I.Ussyukina, A.I.Davydychev. Phys. Lett. B 298 (1993) 363.
- [3] G.t'Hooft, M.Veltman. Nucl.Phys. B153 (1979) 365.
- [4] F.V.Tkachov. Phys.Lett. B100 (1981) 65.

- [5] A.N.Vassiliev, Yu.M.Pis'mak, Yu.R.Honkonen. Teor. Mat.Fiz.(Russ.) 47 (1981) 291;
N.I.Ussyukina. Teor.Mat.Fiz.(Russ.) 54 (1983) 124;
D.I.Kazakov. Teor.Mat.Fiz.(Russ.) 58 (1984) 345.
- [6] N.I.Ussyukina. Preprint Fermilab-Pub-93/241-T;
N.I.Ussyukina. In the X Int.Workshop on High Energy Physics. Zvenigorod, Russia, 1995
(ed. B.B.Levchenko and V.I.Savrin, INP MSU, Moscow, 1996).
- [7] N.I.Ussyukina. Teor.Mat.Fiz.(Russ.) 22 (1975) 300.
- [8] W.N.Baily. Generalized hypergeometric series, University Press, Cambridge, 1935.
- [9] L.Levin. Polylogarithms and associated functions, North Holland, 1981.
- [10] A.V.Stepanova, F.V.Tkachov, in The XII Int. Workshop on High Energy Physics, Samara, Russia, September 1997 [hep-ph/9710242].